

Apollo Medical Centre

(Promoters : Kurnool Hospital Enterprises Ltd.)

43-67/A,N.R. Peta, Kurnool - 518 004, Phone : (08518) 225888, 225889

| | | |
|---|---|---|
| Name : MANOHAR | Bill Date : 06-Jan-2026 8:33 am |  |
| Age : 50 Years | Sample No : 28,28A | |
| Gender : Male | Smpl.Time : 06-Jan-2026 08:37 AM | |
| Bill No : CB26489 | Report Date : 06-Jan-2026 1:09 pm | |
| Ref.Dr. : Dr.RAGHAVENDRA K, MD. DM., | CB26489  | |

DEPARTMENT OF BIOCHEMISTRY

| Test Name | Result | Unit | Bio. Ref. Range | Method |
|--------------------|--------|-------|-----------------------|-----------|
| CREATININE | | | | |
| CREATININE (SERUM) | 1.3 | mg/dL | 0.6 - 1.3 Adult | Enzymatic |
| | | mg/dL | 0.3 - 1.0 Children | |

Creatinine is produced at a fairly constant rate within an individual as a result of breakdown of Creatine within muscle tissue.

Creatinine is freely filtered at the glomerulus and predominantly excreted by the kidneys.

Increased - Old age, glomerulonephritis, pyelonephritis, renal failure, urinary obstruction, CCF, Dehydration, Shock, medicines

like amphotericin B, captopril, cephalosporins etc

Decreased - low muscle mass, females, Malnutrition, Drugs like - Tianoide, Vancomycin etc.,

| | | |
|---|---|--|
| Name : MANOHAR | Bill Date : 06-Jan-2026 8:33 am |  |
| Age : 50 Years | Sample No : 28,28A | |
| Gender : Male | Smpl.Time : 06-Jan-2026 08:37 AM | |
| Bill No : CB26489 | Report Date : 06-Jan-2026 1:09 pm | |
| Ref.Dr. : Dr.RAGHAVENDRA K, MD. DM., | CB26489  | |

DEPARTMENT OF BIOCHEMISTRY

| Test Name | Result | Unit | Bio. Ref. Range | Method |
|---|--------------------|-------|-----------------|-----------|
| GLUCOSE FASTING & PP | | | | |
| GLUCOSE, FASTING , NaF Plasma | 190 | mg/dl | 70 - 100 | GOD - POD |
| URINE SUGAR | NIL | | - | |
| GLUCOSE, POST PRANDIAL (PP), 2 HOURS NAF PLASMA | 333 | mg/dl | 70 - 140 | GOD - POD |
| URINE SUGAR | + + + (THREE PLUS) | | - | |

Comment:

It is recommended that FBS and PPBS should be interpreted with respect to their Biological reference ranges and not with each other.
 Conditions which may lead to lower postprandial glucose levels as compared to fasting glucose levels may be due to reactive hypoglycemia, dietary meal content, duration or timing of sampling after food digestion and absorption, medications such as insulin preparations, sulfonylureas, amylin analogues, or conditions such as overproduction of insulin.

Ref: Marks medical biochemistry and clinical approach

Comment:

As per American Diabetes Guidelines

| Fasting Glucose Values in mg/d L | Interpretation |
|----------------------------------|----------------|
| <100 mg/dL | Normal |
| 100-125 mg/dL | Prediabetes |
| >126 mg/dL | Diabetes |

*** END OF REPORT ***

Printed On : 06-Jan-2026 1:09 pm


Checked By : SAI

CB26489



KINDLY CORRELATE RESULTS WITH CLINICAL FINDINGS & DISCUSS IF NECESSARY.

Page 2 of 2


 Dr.SREELATHA. D
 M.B.B.S, M.D
 Consultant Biochemist

